

[¹⁴C]Pomalidomide and Metabolites in Human Plasma and Urine

Application #AN4240

Conditions

Column: ACE 3 C18
Dimensions: 150 x 4.6 mm
Part Number: ACE-111-1546
Mobile Phase: A: 25 mM ammonium acetate pH 5.5 in H₂O
B: MeOH

Time (mins)	%B
0	0
2	0
38	36
44	100
48	100
50	0



Flow Rate: 0.7 mL/min

Temperature: 30 °C


Detection: Radiometric

Metabolites characterised using LC-MS/MS (positive ion mode)

Plasma (2.5h pooled sample)



Urine (0-72h pooled sample)



1. 3-Aminophthalic acid
2. Hydrolysis product of pomalidomide
3. Hydrolysis product of pomalidomide
4. Glucuronide conjugate of 5-hydroxy pomalidomide
5. Glucuronide conjugate of 5-hydroxy pomalidomide
6. 7-Hydroxy pomalidomide
7. 5-Hydroxy pomalidomide
8. Pomalidomide
9. N-Acetyl pomalidomide

Hoffmann M, Kasserra C, Reyes J, Schafer P, Kosek J, Capone L, Parton A, Kim-Kang H, Surapaneni S, Kumar G. Absorption, Metabolism and Excretion of [¹⁴C]Pomalidomide in Humans following Oral Administration. *Cancer Chemotherapy and Pharmacology* 71, 489-501 (2013) doi 10.1007/s00280-012-2040-6
© 2015 ACT Ltd. All rights reserved. All products are available worldwide.