


### **BIOPHARMACEUTICALS**



## Separation of Amino Acids by HILIC/ Single Quadrupole Mass Detection

359



Representative spectra of 21 standard amino acids separated on a HALO® Glycan column are shown. Amino acid isomers (e.g., Leu, Ile), as well as amino acids: Gln and Lys, are detected under the same mass channels (m/z 132.1 and 147.1, respectively). The elution order for the isomers is Leu first, followed by Ile. The target amino acid is specified with an asterisk (\*) symbol in chromatograms showing multiple peaks (e.g., Cys, Thr, Asn, Glu, Asp).

#### **PEAK IDENTITIES:**

| Amino Acid    | Acronym | Monoisotopic mass <sup>ref 2</sup> [M] | SIR mass <sup>ref 6</sup> [M+H <sup>+</sup> ] |
|---------------|---------|--|---|
| Alanine       | Ala, A  | 89.1                                   | 90  |
| Arginine      | Arg, R  | 174.2                                  | 175   |
| Asparagine    | Asn, N  | 132.1                                  | 133   |
| Aspartic acid | Asp, D  | 133.1                                  | 134   |
| Cysteine      | Cys, C  | 121.2                                  | 122   |
| Cystine ref 7 | n/a     | 240.3 <sup>ref 7</sup>                 | 241   |
| Glutamic acid | Glu, E  | 147.1                                  | 148   |
| Glutamine     | Gln, Q  | 146.2                                  | 147   |
| Glycine       | Gly, G  | 75.1                                   | 76  |
| Histidine     | His, H  | 155.2                                  | 156   |
| Isoleucine    | lle, I  | 131.2                                  | 132   |
| Leucine       | Leu, L  | 131.2                                  | 132   |
| Lysine        | Lys, K  | 146.2                                  | 147   |
| Methionine    | Met, M  | 149.2                                  | 150   |
| Phenylalanine | Phe, F  | 165.2                                  | 166   |
| Proline       | Pro, P  | 115.1                                  | 116   |
| Serine        | Ser, S  | 105.1                                  | 106   |
| Threonine     | Thr, T  | 119.1                                  | 120   |
| Tryptophan    | Trp, W  | 204.2                                  | 205   |
| Tyrosine      | Tyr, Y  | 181.2                                  | 182   |
| Valine        | Val, V  | 117.1                                  | 118   |







### **BIOPHARMACEUTICALS**



# Separation of Amino Acids by HILIC/ Single Quadrupole Mass Detection

### **TEST CONDITIONS:**

Column: HALO 90 Å Glycan, 2.7 µm, 2.1 x 100 mm

Part Number: 92922-605

Mobile Phase A: 85/15 Acetonitrile/ 10 mM ammonium formate, 0.15% Formic Acid

Mobile Phase B: 10 mM ammonium formate pH 3.0, 0.15% Formic Acid

 Gradient:
 Time 0.0
 %B 0.0

 10.0
 5.0

 15.0
 37.5

 15.5
 95.0

 20.0
 95.0

 20.5
 0.0

 25.0
 0.0

Flow Rate: 0.4 mL/min

Column Temperature: 30 °C

Sample Manager Temperature:  $5 \pm 3 \, ^{\circ}\text{C}$ 

Injection Volume: 1 µL

Data Courtesy of: Neshita Kipuw, Pfenex Expression Technology

#### MS CONDITIONS:

**Ionization Mode:** ESI **Polarity:** Positive

Full Scan: m/z 50-1000 Cone Voltage: 15 V

Gas Flow Pressure: 70-100 psi



