




Optimization of Tocochromanols Separation Using Supercritical Fluid Chromatography and HALO® Biphenyl

340



PEAK IDENTITIES

Labeled on chromatogram

TEST CONDITIONS:

Column: HALO 90 Å Biphenyl, 2.7 μ m, 4.6 \times 250 mm

Part Number: 92814-911

Mobile Phase A: CO₂

Mobile Phase B: MeOH

Gradient: Time	%B
0.00	5
5.00	10
6.00	25
12.00	25
13.00	5
14.50	5

Flow Rate: 3.0 mL/min

Back-pressure regulator: 10.0 MPa

Back-pressure: 25.0-31.5 MPa

Temperature: 25 °C

Detection: DAD at 295 nm

Injection Volume: 5.0 μ L

Sample Solvent: 2-propanol

Sample/Solvent ratio: 1:9-3:7, w/v

Tocopherols, tocotrienols, and plastoquinone-8 (PC-8), commonly named tocochromanols/tocols/vitamin E, are fat-soluble antioxidants that play an important role in human health. Applying supercritical fluid chromatography (SFC) and a HALO 90 Å Biphenyl, 2.7 μ m, 4.6 \times 250 mm column for tocochromanols separation the run time can be shortened by three times in comparison to NPLC to 14.5 min. Additionally, the benefit of SFC is that it has green character due to its use of carbon dioxide (CO₂) as the main mobile phase component, which is both non-toxic and readily available. In turn, the HALO® Fused-Core® technology allows for injection of high volume (up to 7-8 μ L) of only diluted plant oils in 2-propanol in a ratio 1:9-3:7, w/v, to the SFC system. The combination of the SFC-DAD and HALO® column technology enabled a rapid and green determination method of nine tocochromanols.

*Data courtesy of Latvian Science Council prepared under project No. lzp-2021/1-0651.



AMT_AN_Rev_0



Made in the USA

INNOVATION YOU CAN TRUST,
PERFORMANCE YOU CAN RELY ON

HALO® and Fused-Core® are registered trademarks of Advanced Materials Technology